

The complete restorative treatment of...



V*aginitis*

50%
of all
women patients



WELLCOME INSTITUTE LIBRARY	
Coll.	welMOMac
Coll.	pam
No.	WP 250
	1949
	S43C

suffer from ***Vaginal Discharge***^①





22500836566

① HARDY, J. A.: "Office Gynecology," J. Missouri M. A. 45:811 (Nov.) 1948.

Complete **Examination**


is necessary to find the infection; many women accept the condition as normal.^②



^② BOEHME, E. J.: "Trichomonas Vaginalis Vaginitis; Diagnosis, Treatment, Causes of Failure in Treatment," S. Clin. North America 25:545 (June) 1945.




normal



A Thick Epithelium


Regardless of causative factor
note the resulting "four hypos."^③

infected • inflamed




Loss of Epithelium


^③ GREENHILL, J. P.: Medical Film,
"Trichomonal and Monilial Vaginitis."



Saturated with Glycogen




An Abundance of Döderlein Bacilli




Sufficient Acidity



Loss of Glycogen



Loss of Döderlein Bacilli



Insufficient Acidity

PH is the index...




Acidity is the most important factor in the treatment of any vaginal infection.^④

^④ KARNAKY, K. J.: "Treatment of Trichomonas Vaginalis," Am. J. Surg. 48:219 (April) 1940.

3

4



Döderlein Bacilli

Thrive 3.8 to 4.4

5

6

7

8




Trichomonas Vaginalis

Thrive 5 to 6



Monilia Albicans


Thrive 5.5 to 6.8



Mixed Infections
(Staph., Strep., B. Coli, etc.)

Thrive 5.8 to 7.8


Floraquin




THE COMPLETE RESTORATIVE TREATMENT...

- Diodoquin
- dextrose
- lactose
- acid

because...




Protozoacide
and fungicide.



... it reestablishes:

{ EPITHELIUM
GLYCOGEN
DÖDERLEIN BACILLI
ACIDITY



Floraquin

EFFECTIVE FOR **ALL** TYPES OF VAGINITIS



Trichomonas Vaginalis




Monilia Albicans




Mixed Infections




Senile Vaginitis



Floraquin regimen




POWDER



Office Treatment

Watch progress and maintain cooperation of patients.




TABLETS



Home Treatment

Continue through at least two menstrual periods.




Suggested Douche—

four tablespoonfuls (2 oz.) of white vinegar (or 5 per cent acetic acid) to 2 quarts of warm water.

Home Treatment


Supplement with acid douches for hygienic purposes.

Maintain Acidity



Prevent Reinfection

Use of Diodoquin tablets orally as adjunctive treatment if reinfection from *Trichomonas intestinalis* is suspected.





SEARLE

RESEARCH IN THE SERVICE OF MEDICINE